



THE REPUBLIC OF UGANDA
MINISTRY OF HEALTH


NATIONAL TREATMENT ALGORITHMS FOR SEXUALLY TRANSMITTED DISEASES IN UGANDA

December 2003


Prepared by:
**STD Control Unit
STD/AIDS Control Programme
Ministry of Health:**

**AIM District Programme
Uganda**


MANAGEMENT OF URETHRAL DISCHARGE




MANAGEMENT OF GENITAL ULCER DISEASE




MANAGEMENT OF VAGINAL DISCHARGE SYNDROME




MANAGEMENT OF LOWER ABDOMINAL PAIN IN WOMEN




MANAGEMENT OF INGUINAL BUBO



MANAGEMENT OF ACUTE SCROTAL SWELLING



MANAGEMENT OF BALANITIS IN MEN



MANAGEMENT (OPHTHALMIA NEONATORUM):

Conjunctivitis with pus in the eyes of the newborn

**Clean the eye with saline or clean water,
Always wear gloves
Cover the inflamed eye with gauze before you open the eyelids
for your own protection.
Plus
Ceftriaxone 50mg / Kg body weight (Max of 125mg) i.m. stat or
Kanamycin 75mg stat
Plus
Erythromycin syrup 15 mg/kg body weight 6 hourly for 14 days**

-
- Treat mother and father for cervical and urethral discharge respectively.
 - Prophylaxis for ophthalmia neonatorum: Tetracycline eye ointment stat or Silver nitrate 1% stat, immediately after birth.